

Effects of aqueous extracts of *Azadirachta indica* (neem) on the larvae of *Culex pipiens* L. (Diptera: Culicidae)

By

**Ammar Ahmed Sultan AL-qeraquoly and Ghassan Hamdan Jamel AL-kaisy
University of Diala
Collage of Education
Biology Department**

ABSTRACT

The effect of crude aqueous extracts of *Azadirachta indica* (neem) against the larvae of *Culex pipiens* L. was investigated. Exposure of the larvae to undiluted extracts of seed oil, leaf and bark for 12 hours led to 100, 98, and 48% mortality, respectively. Dilution of these extracts also resulted in mortality of the larvae. We suggest that the seed oil and leaf extract of neem contain properties that could be developed and used in control of *Culex* in Iraq

INTRODUCTION

***Azadirachta indica* A. Juss (neem) is a member of the family meliaceae in the order Fagales of the Dicotyledonous class in the plant kingdom. When fully grown, it can reach 24m in height with dense wide spreading crown. The crown bears large number of compound leaves each with 5-9 pairs of leaflets. Flowering is twice a year (April/May and August/September). When fertilized one seeded fruits are produced with fibrous cover. According to Randwanski(1977), a tree can yield between 20-50 kg of fruits annually.**

The neem has been reported to contain several biologically active constituents such as azadirachtin (Naganishi, 1975), meliantriol (Lavie et al., 1965), salanin (Warthen et al., 1978), as well as nimbin and nimbidin (Shin-foon, 1984). Garg and Bhakuni (1984) reported salanolide (a meliacin) as one of the bitter principles in neem seed oil. Furthermore, Raman and Santhanagopalan (1979) reported tignic acid (5-methyl-2-butanoic acid) as part of the seed constituents. This compound is believed to be responsible for the distinctive odour of

the neem oil. The limonoids are freely soluble in organic solvents such as hydrocarbons, alcohol and ketones, but are sparingly soluble in water.

Azadirachtin, a limonoid, has been reported to have adverse effect on endocrine system of a bean beetle, *Epilachna varivestis*, and cause sterility in the female insects (Schmuthere et al., 1981). Schluter and Schulz (1983) also reported this compound to cause degradation in larvae from molting. Insect larvae treated with meliantriol concentration as low as 0.2 mg/larvae were observed to cease to feed. In recognition of these important properties of neem on insect. This preliminary effort is made to test the effect of crude extract of neem seed oil, leaf and root on the larvae of *Culex pipiens* L. that is responsible for transmitting the malarial parasite, Plasmodium sp. The choice of crude extract was to ensure adaptability by the local populace.

MATERIALS AND METHODS

Extracts of seed kernel, leaf and bark of neem were prepared from plant material. Extraction of all the parts was carried out using local methods. This is allowed for easy adoption by the local population. Seed kernel extract was prepared from sundried seeds that were pounded in a wooden mortar into a paste. The paste was compressed to extract oil which was collected in a round bottom conical flask. To prepare the leaf extracts, fresh leaves were collected, washed and crushed in a motor into a watery paste. This was then gradually passed on wire gauze on top of a conical flask and carefully pressed to extract the liquid which collected in a flask placed below the wire gauze. The flask containing the liquid was then sealed for further use. Bark extract was prepared from fresh pieces of stem bark removed from a full-grown neem tree. The pieces were dried, powdered and sieved to separate finer particles from granules and fibres. From the finer material, 10 g was measured and dissolved in 100 ml distilled water in a conical flask.

Culex larvae were collected in a plastic bucket from a pool of stagnant water. The bucket containing the larvae was kept in a netted enclosure in the laboratory at room temperature ($30\pm 2^\circ\text{C}$).

Three sets of four petri dishes were laden with moist filter paper and placed on a laboratory bench. Using a metal loop, 20 *Culex*

larvae were transferred from the plastic bucket into each of the twelve petri dishes. The sets of petri dishes were labeled a,b,c and d. In the first set, 20 mls of the seed oil extract were added into dish (a), 10 mls in (b), and 5 mls in (c). In petri dish (d), 20 mls of distilled water was added as control. In the second and third sets of petri dishes, leaf and bark extracts were added in the same manner as for seed oil in the first set. The effects of the three extracts were monitored by counting the number of dead larvae at one-hour interval for 12 h. three replicates of the experiments was conducted and mean taken. The data obtained was statistically assessed to determine variance and standard error of the mean.

RESULTS AND DISCUSSION

The effects of the plant extracts on *Culex* are presented in Tables 1,2 and 3. When 20 ml of the seed oil extract was applied, all the larvae died within 12 h. Decreasing the amount of the seed oil extract to 10 and 5 ml, reduced the mortality to 98 and 85%, respectively, after 12 h. For the leaf extract (Table 2), 83% mortality was recorded when the larvae were treated with 20 ml extract, while 75 and 68% mortality was recorded with 10 and 5 ml extracts, respectively, after 12 h. The highest mortality for the bark extract is 48% with 20 ml.

This work demonstrates the potency of *A. indica* in the control of *Culex* larvae. Seed oil appeared as the most lethal among the various parts tested. The high mortality recorded for seed oil extract might be attributed to deficiency of dissolved oxygen in the water. Previous studies have shown that extracts of some plant parts do possess insecticidal effects. This result compare favourably with that from other species, for example *Ambrosia meritima* require application of 1000 ppm to achieve result similar to that of leaf extract (Sharif and El-Sawy 1962). Extracts of *vernonia amygdlina* leaves can cause about 70% mortality in *Streptococcus marcescens* and *S . pyogenes* (Dangoggo et al., 2002).

Table 3. Mortality rate of different quantity of neem bark extract on *Culex pipiens* L. larvae

Bark extract Concentration (MI)	Time (hours)													Total	%	Mean	SE (±)
	1	2	3	4	5	6	7	8	9	10	11	12					
20	4	2	1	1	3	2	2	5	4	4	1	0	29	48	2.42	1.56	
10	0	0	3	1	0	4	2	0	2	0	1	1	14	23	1.17	1.33	
5	0	0	0	3	1	2	2	0	2	0	1	1	12	20	1.0	1.04	
control	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0	

REFERENCES

- Dangoggo SM, Faruq UZ, Manga SB (2002). Antibacterial Assessment and phytochemical screening of *Vernonia amygdalina* leaves. *Nig. J . Basic appl. Sci.* 11:1-8.
- Garg HS, Bhakuni DS (1984). *Phytochem.* 23:2383. Cited in MM Iwu (1993). *Hand book of African medicinal plants.* CRC press, London pp. 124-128.
- Lavie D, Jain MK, Shpan-Gabrielith SR (1965). A locust phagorepellent from two *Melia* species. *Chem. Commun.* 910-1911.
- Naganishi K (1975). Structure of the insect antifeedant azadirachtin. In: *Recent advances in phytochemistry*, VC Runeckles (ed.), plenum, New York, N.Y. Vol.5 pp. 283-298.

- Radwanski, S (1977). Neem tree 2. Uses and potential uses. World crops and Livestock. July/August, pp. 167-168.

- Raman H, Santhanagopalan S (1979). Indian J.chem. Sect. Borg. Ind. Med. Chem. 17: 169. cited in MM Iwu (1993). Hand book of African medicinal plants. CRC press, London pp. 124-128.

- Schluter U, Schulz WD (1983). Structural damages caused by neem in *Epilachna verivestis*: a summary of histological and ultrastructural damage. 1. Tissues affected in larvae. Proc. 2nd Int. neem conf. Rottach. Egern, p. 294.

- Schmutterer H, Ascher KRS, Rembold H (1981). Natural pesticides from the neem tree. Proc. 1st Int. neem Conf. Rottach-Egem, p 294.

- Sherif, AF, El Sawy MFED (1962). Using damissia in field trials as a control method of snails vector. In first symposium of bilhaziasis. Govt. printing office, Cairo. P 1138.

- Shin- Foon C (1984). The active principles and insecticidal properties of some chinese plants, with special reference to Meliaceae. Pp. 255-262.

- Warthen JD Jr, Uebel EC, Dutky SR, Lusby WR, Finegold H (1978). An adult housfly feeding deterrent from neem seeds. UD Agric., Agric. Res. Results RR-NE 2.