

## Examination of Catalase enzyme in fresh leaves and study some characters of it

\* A.K.AL-Anbari

### Summary

#### CATALASEIEC:

1.1.1.6) was eliminated in the lettuce *Lactuca sativa* ,cabbage *Brassia oleracea*, spinach *Spiacia olearacea*, Chard *Beta vulgarer* leaves , enzyme assay ,protein contains , optimum temperature of .

Activity & stability ,optimum ph of activity & stability .

The result shower The chard *Beta bulgaria* leaves was biggest activity of database enzyme was (256.897) U/mg , protein contain was ( 4.797) mg/mi , optimum temperature of activity 50c was(0.256) U/ml , optimum temperature of stability 50C to chelation 30 min , CAT has optimum ph of activity 8 was (0.608) U/ml , optimum ph of stability was (1.455)U/ml

#### Information

Catalase plays a major role in the protection of tissues from the toxic effect of  $H_2O_2$  . CAT also removes electrons that can lead to the production of  $O_2$  free- radical ( Abassi et al ; 1998) . CAT has been found in all aerobic cells containing cytochrom & CAT enzyme is one of the earliest enzyme had been studied & purified ( Percy ; 1984 ). It is primarily an intra cellular enzyme . The enzyme usually it have been sensitive from extreme temperature degree , the extreme temperature cause changes in primary secondary & active structure of protein & denaturation of it as well as the changes of PH effects in natural of enzyme . The extreme ph decreased activity & \* Biology department collage of education / university of dyila

stability of enzyme . Optimum temperature of activity & ph depended on the source of enzyme ( Aydemir et al ; 2003) The aim of the present study was to characterize the CAT from the local varieties lettuce *Lactuca sativa* , cabbage *Brassia oleracea* , spinach *Spiacia olearacea*, chard *Beta vulgarer* leaves & enzyme assays , protein contains , Optimum temperature and ph of activity & stability.

#### Materials & methods

##### Chemicals

Hydrogen peroxide , Phosphate buffer (  $Na_2 HPO_4$  , $K_2HPO_4$  ) , Bovine

serum albumin (BSA) ,Ammonium collide , Acitate buffer , iris buffer.  
Enzyme assay CAT activity was determined of fresh leaves at 25c according to A1 - Alwani ( Alwani ;2006 ) The reaction mixture contained (10 mM) h2O2 in a (60mM) sodium potassium phosphate buffer ph 7.4 & 20 ul crude than addition (34.2M) ammonium molybdate .CAT activity was estimated by decreased in absorbency of h2O2 at 504nm .according to equation : -

$$\text{Catalase Activity} = \frac{\text{Sample - Blank 1}}{\text{Blank 2 - Blank 3}} \times 271$$

if Blank 1= 1ml (H2O2 + sodium potassium phosphate buffer) + 1 ml (ammonium Molybdate ) + 0.2 sample .

Blank 2= 1ml (H2O2 + sodium potassium phosphate buffer) + 1ml (ammonium Molybdate ) + 0.2 ( sodium potassium phosphate buffer) .

Blank 3| 1ml ( sodium potassium phosphate buffer) + 1ml (ammonium Molybdate ) + 0.2 sodium potassium phosphate duffer) .

Determination of protein Protein amount for CAT was done according to method of Mohamed & abd all (1996 , محمد و عبدالله)with bovine serum albumin as standard

Effect of ph

The effect of ph of CAT activity was investigated in the range of ph 4-5 (0.05M) acetate buffer , ph 6-8 (0.05M) phosphate buffer , ph 9-10 (0.05M) tries buffer .Then 10 ul enzyme solution was mixed with 0.99 ml buffers at various ph values incubated at 25c for 1 h . Thereafter retained enzyme activity was measured as indieatedin Aebi (Aebi ;1984 ) after that optimum ph was determined .

Effect of temperature The effect of temperature on CAT activity was investigated by incubating the enzyme solution in ( 0.05 M ) phosphate buffer ph 7.0 for 30 min in the range of 10 -70 C. Then 10 ul enzyme solution was mixed with 0.99 ml phosphate buffer & incubated , Thereafter retained enzyme activity was measured by absorption at 240 nm according to Aebi (Aebi ; 1984 ) after that optimum temperature was determined .

Optimum ph to stability To CAT stability was determined by incubation of enzyme solution with previously to prepare buffers at 1:1 at 25c for 60 min & measured by absorption at 240 nm .

Optimum Temperature to stability Various temperatures were used to CAT stability was determined by incubation of enzyme solution at 50 C at seven different time ( 10 - 70 min in bath water , Than 20 ul enzyme solution mixed with 0.99 ml h2O2 (65 mM) & Sodium phosphate buffer ( 60 mM) ph 7.0 to stopped the enzyme work & measured by absorption at 240 nm .

**Results & discussion**


(Table 1 ) shown the biggest CAT total activity was determined at chard leaves were 6161.68 units , that affected to contain of protein was measured to be (4.797) mg/ml

Table (1) CAT activity & protein

Crude extract Total activity Units/mg) ( mg / ml)	Specific activity (Units)	Protein ( Units/ mg protein)	Activity ( mg/ml )	Total protein ( mg/ml )
L.sativa 732.17	101.409	1.444	146.435	507.040
B.oleracea 784.473	81.124	1.934	156.897	405.622
S.olearacea 4716.350	197.502	4.776	943.270	987.510
B.vulgaris 6161.684	256.897	4.797	1232.336	284.486

The CAT activity shown in (Fig 1) increasing when the incubating temperature until it was (0.256)U/ml at 50C, Then the activity was


decreasing because the increasing temperature increased the speed of reaction due to increasing movement energy to enzyme molecules ( Al-Akarky ; 2002)



(Fig 1) effect of Temperature in enzyme activity(CAT)


The thermal stability appear in (fig 4 ) CAT stability in the enzyme solution kept storge there activity when incubating 50C° 30 min was (0.604)U/ml

The optimum temperature we determined was similar to that given in previous studies ( Demir ; 2004)




(Fig4) effect of Temperature in stability of enzyme (CAT)

In the study of effect at different pH of CAT activity shown in (fig 3 ) the biggest activity was (0.608)U/ml at pH 8 ,Then decreasing in crude cause the denaturation of protein molecules , however the same results were arrived ( Yoruk ; 2005 ;Gholamhoseinian ; 2006 )



(Fig 2)effect Of pH in to enzyme activity(CAT)

Enzyme stability was expressed as remaining activity ,The stability at different pH values was shown (fig 4 ) when CAT was stored at pH 8 was (1.455)U /ml other studies has been reposed exhibit sharp optimum pH ( Torres ; 2003)



( Fig 4 ) effect of PH in stability of enzyme

التحري عن انزيم الكاتليز في الأوراق الخضراء ودراسة بعض خصائصه

\*أسيل كاظم الانباري

### المستخلص

تم التحري عن انزيم الكاتليز CAT في أوراق الخس *Lactuca sativa* واللهاة *Brassia oleracea* والسبانخ *Spiacia oleracea* والسلق *Beta vulgaris* وتم تقدير الانزيمية له اذ أظهرت أوراق السلق أعلى فعالية بلغت (256.897)  $\mu\text{g}$  وتمت دراسة محتوى الاوراق من البروتين الكلي الذي تفوقت اوراق السلق والسبانخ في اعلى محتوى بلغ ( 4.776  $\text{ml}\backslash\text{mg}$ (4.797 على التوالي كما تم اختيار لدرجة الحرارة المثلى لفاعلية وثبات الانزيم والذالة الحامضية المثلى لفاعلية وثبات عمل الانزيم في اوراق السلق اظهرت الدراسة النتائج التالية:.

كانت درجة الحرارة المثلى لفاعلية الانزيم 50 C بلغت عندها فاعليته (0.256)  $\mu\text{ml}$  و اظهر الانزيم ثباتا عند حضنه لمدة 30 دقيقة في درجة 50 م وصل ( 0.604)  $\mu\text{ml}$  اما الذالة الحامضية المثلى للفاعلية كانت عند ( 8 ) (ph) وبلغت (0.608)  $\mu\text{ml}$  وابدى الانزيم ثباتا عند نفس الذالة الحامضية ووصل الى (1.455)  $\mu\text{ml}$

References

- محمد عبد العظيم كاظم وليلى نجم عبد الله .فسلجة النبات العلمي .وزارة التعليم العالي والبحث العلمي .جامعة بابل (1996).
- Abassi , N.A.,M.Kushad & Endress .Active oxygen scavenging enzymes activities in developing apple flowers & fruits . Sientia Horticult - urac . 74 : 183 - 194 .(1998)
- Aebi , H.catalase In :unpacker (Edl,methods in enzymology . Academic press .orlando . 105 : 121 - 126 .( 1984).
- AL-Karky , Layla Othman . Extraction , purification & unease in seeds of wheat & study of characterization of effect of some inhibitions .thesis of master College of Science for Women University of Baghdad .(2002).
- AL-Alwani , Basheer Abdul Hamzah agging causes intrums of oxidative hypothesis in relation to adventitious roots formation in mung bean cutting .thesis of doctor college of scinence University of Babylon (2006) .
- Aydemir , T. & K. Kuru . Purification & partial characterization of catalyst from chicken erythrocytes & effect of various in hibhtors on enzy- me activity Turk .J.Chem . 27 : 85 - 97 .(2003) .
- Demir Hpurification & characterization of polyphenol oxidate from the Ercic apple . Bulletin of pure & applied sciences chemistry . 42:1-4 ( 2004) .
- Gholamhoseinian ,A.; N. Ghaemi & F. Rahimi .partial PUification & properties of database from Bssia oleracea .Asian .J.of plant sciences 5 (5) : 827- 831 .(2006) .
- Percy ,M. E. Catalase an old enzyme with anew role ? Can. J.Cell . Biol 62: 1006 - 1014 . ( 1984 )
- Torres ,R. M. C. Valentines ,J. Vsall & Vinas . Possible involvement of hydrogen peroxide in development of resistance mechanisms in golden delicious apple fruit . postharvest. Biology & Technology . 27: 235 - 242 . (2003)



- Yoruk, i H. Demir ,K. Ekici ,& A. Savran purification & properties of catalyst from Van apple ( Golden Delicious ) .  
Pakistan .J.of Nutri - lion . 4 (1) : 8 - 10 . (2005) .